


Scarlet Tanager

Virginia's Second Breeding Bird Atlas


Environmental Associations




Scarlet Tanager © Julie Davis;
Cornell Lab of Ornithology |
Macaulay Library




Percent Agricultural Cover



Percent Shrubland/Grassland
Cover



Percent Forest Cover



Size of Largest Forest Patch

The graphs presented on this page illustrate the probability that a species will occur within a block based on the values of significant environmental variables for that block in the Second Atlas species' distribution models. Probability of occurrence ranges from 0 to 1 (vertical axis). Depending on the graph, the horizontal axis depicts 1) the percentage of a block covered by a habitat type (e.g., grassland and shrubland) or 2) a value from low to high for climate variables (e.g., mean annual rainfall) and landscape-level features (e.g., diversity of habitat types) within a block. The dark line indicates the relationship between occurrence and the variable. There is a 95% probability (confidence interval) that this relationship falls between the gray lines. The black tick marks at the bottom of the graph show where individual blocks fall on the horizontal axis, to demonstrate the spread of values for the blocks in the model.


Scarlet Tanager

Virginia's Second Breeding Bird Atlas


Environmental Associations




Scarlet Tanager © Julie Davis;
Cornell Lab of Ornithology |
Macaulay Library



Diversity of Habitat Types



Variation in Annual Rainfall



Mean Annual Rainfall

The graphs presented on this page illustrate the probability that a species will occur within a block based on the values of significant environmental variables for that block in the Second Atlas species' distribution models. Probability of occurrence ranges from 0 to 1 (vertical axis). Depending on the graph, the horizontal axis depicts 1) the percentage of a block covered by a habitat type (e.g., grassland and shrubland) or 2) a value from low to high for climate variables (e.g., mean annual rainfall) and landscape-level features (e.g., diversity of habitat types) within a block. The dark line indicates the relationship between occurrence and the variable. There is a 95% probability (confidence interval) that this relationship falls between the gray lines. The black tick marks at the bottom of the graph show where individual blocks fall on the horizontal axis, to demonstrate the spread of values for the blocks in the model.